

2003 Health and Preventive Practices Status Report

from Missouri County-level Study



Stoddard County**

	County Prevalence (%)*	Southeast Region Prevalence (%)*	State Prevalence (%)*
G eneral			
Health Status (Fair or Poor)	22.7	23.2 H	16.9
No Health Coverage	20.3	17.5 H	12.8
B ehavioral Risk Factors			
Physical Inactivity	34.5 H	30.8 H	24.0
Current Smoker	39.4 H	28.1	26.4
Overweight (BMI 25-29.9)	34.8	36.9	35.3
Obese (BMI ≥ 30)	26.6	26.9	23.4
D isease and Conditions			
High Blood Pressure	33.7	30.1	28.9
High Cholesterol (35 years or older)	46.3 H	38.2	37.1
Asthma	15.8	12.3	10.8
Diabetes	6.4	7.5	7.2
Activity Limitation	18.1	20.6	17.9
C ancer Screening			
<i>Women</i>			
Never had a Mammogram	15.4	20.5 H	13.4
No Mammogram or Clinical Breast Exam in Last Year:			
40-49 years of age	42.6	58.0	49.4
50-64 years of age	56.6	44.8	35.9
65 and older	44.2	52.9	47.4
Never had a Pap Test:			
18-69 years of age	1.7	3.8	5.1
70 and older	11.0	8.2	6.4
No Pap Test in Last Year (18-29)	11.1	23.5	25.8
No Pap Test in Last 3 Years (30-69)	20.6	15.4	13.8
<i>Men and Women 50 Years or Older</i>			
Blood Stool Test:			
Never had a Blood Stool Test	62.8	63.1 H	57.0
No Blood Stool Test in Last Year	87.7 H	83.7 H	77.5
Sigmoidoscopy/Colonoscopy			
Never had a Sigmoidoscopy or Colonoscopy	52.0	59.0 H	49.2
No Sigmoidoscopy or Colonoscopy in Past Five Years	64.3	67.7 H	57.0

*Prevalence rates are age adjusted to the 2000 U.S. standard population

■ indicates that the prevalence rate is not significantly above or below state prevalence

H indicates that the prevalence rate is significantly above the state prevalence

L indicates that the prevalence rate is significantly below the state prevalence

Unless otherwise noted respondents are 18 years of age or older

** The best indicators for Stoddard county were obtained by combining data from Stoddard and New Madrid counties.